

e, E. Sundquist, and G. Winston

nautics and stration

ce Flight Center yland 20771

The NASA STI Program Office ... in Profile

Since its founding, NASA has been dedicated to the advancement of aeronautics and space science. The NASA Scientific and Technical Information (STI) Program Office plays a key part in helping NASA maintain this important role.

The NASA STI Program Office is operated by Langley Research Center, the lead center for NASA's scientific and technical information. The NASA STI Program Office provides access to the NASA STI Database, the largest collection of aeronautical and space science STI in the world. The Program Office is also NASA's institutional mechanism for disseminating the results of its research and development activities. These results are published by NASA in the NASA STI Report Series, which includes the following report types:

- TECHNICAL PUBLICATION. Reports of completed research or a major significant phase of research that present the results of NASA programs and include extensive data or theoretical analysis. Includes compilations of significant scientific and technical data and information deemed to be of continuing reference value. NASA's counterpart of peer-reviewed formal professional papers but has less stringent limitations on manuscript length and extent of graphic presentations.
- TECHNICAL MEMORANDUM. Scientific and technical findings that are preliminary or of specialized interest, e.g., quick release reports, working papers, and bibliographies that contain minimal annotation. Does not contain extensive analysis.
- CONTRACTOR REPORT. Scientific and technical findings by NASA-sponsored contractors and grantees.

- CONFERENCE PUBLICATION. Collected papers from scientific and technical conferences, symposia, seminars, or other meetings sponsored or cosponsored by NASA.
- SPECIAL PUBLICATION. Scientific, technical, or historical information from NASA programs, projects, and mission, often concerned with subjects having substantial public interest.
- TECHNICAL TRANSLATION.
 English-language translations of foreign scientific and technical material pertinent to NASA's mission.

Specialized services that complement the STI Program Office's diverse offerings include creating custom thesauri, building customized databases, organizing and publishing research results . . . even providing videos.

For more information about the NASA STI Program Office, see the following:

- Access the NASA STI Program Home Page at http://www.sti.nasa.gov/STI-homepage.html
- E-mail your question via the Internet to help@sti.nasa.gov
- Fax your question to the NASA Access Help Desk at (301) 621-0134
- Telephone the NASA Access Help Desk at (301) 621-0390
- Write to: NASA Access Help Desk

NASA Access Help Desk NASA Center for AeroSpace Information 7121 Standard Drive Hanover, MD 21076-1320

NASA/TM—2000–209891, Vol. 245



Technical Report Series on the Boreal Ecosystem-Atmosphere Study (BOREAS)

Forrest G. Hall and Sara K. Conrad, Editors

Volume 245 BOREAS TGB-12 ²²²Rn Activity Data over the NSA

Susan Trumbore, University of California, Irvine Eric Sundquist and Greg Winston, U.S. Geological Survey, Woods Hole, Massachusetts

National Aeronautics and Space Administration

Goddard Space Flight Center Greenbelt, Maryland 20771

	A!1-1-1 C	
	Available from:	
NASA Center for AeroSpace Information		National Technical Information Service
7101 G. 1 1D.		
7121 Standard Drive		5285 Port Royal Road
Hanover, MD 21076-1320		Springfield, VA 22161
Price Code: A17		Price Code: A10
THE COUE, AT		rnce Coue. A10

BOREAS TGB-12 222Rn Activity Data over the NSA

Susan Trumbore, Eric Sundquist, Greg Winston

Summary

The BOREAS TGB-12 team made measurements of soil carbon inventories, carbon concentration in soil gases, and rates of soil respiration at several sites to estimate the rates of carbon accumulation and turnover in each of the major vegetation types. Sampling strategies for soils were designed to take advantage of local fire chronosequences, so that the accumulation of carbon in regrowing mosses could be determined. All the data are used to (1) calculate the inventory of carbon and nitrogen in moss and mineral soil layers at NSA sites, (2) determine the rates of input and turnover (using both accumulation since the last stand-killing fire and radiocarbon data), and (3) link changes in soil respiration rate to shifts in the ¹⁴C content of soil CO₂ to determine the average 'age' respired CO₂. These ²²²Rn activity data were collected from 15-Nov-1993 to 16-Aug-1994 over the NSA sites. They are useful for determining the rate of gas exchange between soil and the overlying atmosphere. The data in this data set are stored in tabular ASCII files.

Table of Contents

- 1) Data Set Overview
- 2) Investigator(s)
- 3) Theory of Measurements
- 4) Equipment
- 5) Data Acquisition Methods
- 6) Observations
- 7) Data Description
- 8) Data Organization
- 9) Data Manipulations
- 10) Errors
- 11) Notes
- 12) Application of the Data Set
- 13) Future Modifications and Plans
- 14) Software
- 15) Data Access
- 16) Output Products and Availability
- 17) References
- 18) Glossary of Terms
- 19) List of Acronyms
- 20) Document Information

1. Data Set Overview

1.1 Data Set Identification

BOREAS TGB-12 222Rn Activity Data over the NSA

1.2 Data Set Introduction

This data set contains seasonal variations in the ²²²Rn soil gas concentrations. The data presented represent sampling from the fall of 1993 to the summer of 1994.

1.3 Objective/Purpose

The objective of these data is to estimate rates of carbon input, turnover, and accumulation in the soils of each of the major vegetation types at the BOReal Ecosystem-Atmosphere Study (BOREAS) study sites. In addition, the data will be used to relate the estimates of soil carbon dynamics to ecosystem models of the carbon cycle, other measures of carbon cycling dynamics, regional models of soil carbon accumulation, and spatial and temporal models of soil moisture and drainage.

1.4 Summary of Parameters

²²²Rn activity and soil depth were measured over the BOREAS Northern Study Area (NSA). Other parameters measured simultaneously include CO₂ concentrations in the soil atmosphere, reported in a separate data set. We concentrated on making measurements during the winter months. Sites for measurements were those also used by Trace Gas Biogeochemistry (TGB)-01 (Crill) and TGB-03 (Moore) for flux measurements and concentration profiles during the growing season.

1.5 Discussion

Radon data may be used to estimate the effective diffusivity of gases within the soil atmosphere, or to test models that estimate effective diffusivity from parameters like air-filled porosity. Details may be found in Davidson and Trumbore (1995).

1.6 Related Data Sets

BOREAS TGB-12 222Rn Flux Data over the NSA

BOREAS TGB-12 Isotopic Carbon Dioxide Data over the NSA

BOREAS TGB-05 Fire History of Manitoba 1980 to 1991 in Raster Format

BOREAS TE-18 Landsat TM Physical Classification Image of the NSA

BOREAS AFM-12 1-km AVHRR Seasonal Land Cover Classification

BOREAS Regional Soils Data in Raster Format and AEAC Projection

BOREAS Soils Data over the SSA in Raster Format and AEAC Projection

BOREAS TGB-01 Soil CH4 and CO2 Profile Data over the NSA

2. Investigator(s)

2.1 Investigator(s) Name and Title

Susan Trumbore Department of Earth System Science UC Irvine

Jennifer Harden U.S. Geological Survey Menlo Park, CA

Eric Sundquist U.S. Geological Survey Woods Hole, MA

Eric Davidson Woods Hole Research Center Woods Hole, MA

2.2 Title of Investigation

Input, Accumulation and Turnover of Carbon in BOREAS NSA Soils (TGB-12)

2.3 Contact Information

Contact 1:

Susan Trumbore
Department of Earth System Science
University of California
Irvine, CA 92697-3100
(714) 824-6142
(714) 824-3256 (fax)
setrumbo@uci.edu

Contact 2:

Greg Winston
U.S. Geological Survey
Quisett Campus
Woods Hole, MA 02543
Winston@nobska.wr.usgs.gov

Contact 3:

Eric Sundquist U.S. Geological Survey Quisett Campus Woods Hole, MA 02543 (508) 457-2397 sundquist@nobska.wr.usgs.gov

Contact 4:

Jeffrey A. Newcomer Raytheon ITSS Code 923 NASA GSFC Greenbelt, MD 20771 (301) 286-7858 (301) 286-0239 (fax) Jeffrey.Newcomer@gsfc.nasa.gov

3. Theory of Measurements

A static chamber method was used to determine radon fluxes. Chambers are described in Winston et al. (1997) and problems with chamber sampling in Stephens and Sundquist (1998). After chamber emplacement, samples of chamber air were taken at intervals of 30 to 45 minutes and used to fill pre-evacuated Lucas counting cells. The flux is determined from the slope of radon activity versus time. Special pits were instrumented with thermistors (for monitoring soil temperature), Time Domain Reflectometry (TDR) probes (for monitoring soil water content), and soil gas probes (1/8" stainless steel tubing, perforated at one end and inserted 50 to 100 cm laterally into the soil pit wall, capped with 1/8" swagelock union fittings sealed with a septum). Further details are given in Winston et al. (1997) and in Section 4, below.

²²²Rn gas samples were obtained using gas-tight syringes (60 cc), dried (by passing through drierite), and then allowed to fill an evacuated Lucas counting cell coated with phosphor scintillant and decay counted. A description of radon sampling and analysis procedures is given in Davidson and Trumbore (1995).

4. Equipment

4.1 Sensor/Instrument Description

Because all of the equipment used in this project is common to many other projects and no special procedures were used, description detail has been minimized in this section, and the reader is referred to the appropriate publications.

- Davidson and Trumbore, 1995
- Mathieu et al., 1988
- Stephens and Sundquist, 1998
- Winston et al., 1997

Radon measurements were made using Pylon alpha scintillation counters.

4.1.1 Collection Environment

Samples were collected under all environmental conditions. Most measurements represent winter conditions.

4.1.2 Source/Platform

Soil.

4.1.3 Source/Platform Mission Objectives

The objective was to determine the soil ²²²Rn activity.

4.1.4 Key Variables

The key variables measured were ²²²Rn activity and soil depth. ²²²Rn fluxes are also reported.

4.1.5 Principles of Operation

None given.

4.1.6 Sensor/Instrument Measurement Geometry

Not applicable.

4.1.7 Manufacturer of Sensor/Instrument

None given.

4.2 Calibration

Radon cell efficiencies were determined in the lab using background air and 226Ra of known activity absorbed to manganese fibers and sealed in a tube. Counting cell backgrounds were determined using ambient air (see Davidson and Trumbore, 1995, for details of radon measurement methods).

4.2.1 Specifications

None given.

4.2.1.1 Tolerance

None given.

4.2.2 Frequency of Calibration

None given.

4.2.3 Other Calibration Information

None.

5. Data Acquisition Methods

None given.

6. Observations

6.1 Data Notes

None given.

6.2 Field Notes

None given.

7. Data Description

7.1 Spatial Characteristics

7.1.1 Spatial Coverage

The North American Datum of 1983 (NAD83) coordinates of the sites were:

NSA-Old Black Spruce (OBS):	55.88007N,	98.48139W
NSA-Old Jack Pine (OJP):	55.92842N,	98.62396W
NSA-Young Jack Pine (YJP):	55.89575N,	98.28706W
NSA-Gillam Road:	55.9055 N,	97.70872W

7.1.2 Spatial Coverage Map

Not available.

7.1.3 Spatial Resolution

None given.

7.1.4 Projection

Not applicable.

7.1.5 Grid Description

Not applicable.

7.2 Temporal Characteristics

7.2.1 Temporal Coverage

Soil gas flux and concentration measurements were made from November 1993 to August 1994.

7.2.2 Temporal Coverage Map

Not applicable.

7.2.3 Temporal Resolution

The temporal resolution of the measurements was variable. Some sites were visited once and other were visited multiple times.

7.3 Data Characteristics

7.3.1 Parameter/Variable

The parameters contained in the data files on the CD-ROM are:

7.3.2 Variable Description/Definition

The descriptions of the parameters contained in the data files on the CD-ROM are

Column Name	Description
SITE_NAME	The identifier assigned to the site by BOREAS, in the format SSS-TTT-CCCCC, where SSS identifies the portion of the study area: NSA, SSA, REG, TRN, and TTT identifies the cover type for the site, 999 if unknown, and CCCCC is the identifier for site, exactly what it means will vary with site type.
SUB_SITE	The identifier assigned to the sub-site by BOREAS, in the format GGGGG-IIIII, where GGGGG is the group associated with the sub-site instrument, e.g. HYD06 or STAFF, and IIIII is the identifier for sub-site, often this will refer to an instrument.
DATE OBS	The date on which the data were collected.
SOIL_DEPTH	The depth below the soil surface at which the measurement was taken.
RADON222_ACTIVITY	RADON-222 activity in air or soil air space. (A becquerel is 1 decay per second.)
CRTFCN_CODE	The BOREAS certification level of the data. Examples are CPI (Checked by PI), CGR (Certified by Group), PRE (Preliminary), and CPI-??? (CPI but questionable).
REVISION_DATE	The most recent date when the information in the referenced data base table record was revised.

7.3.3 Unit of Measurement

The measurement units for the parameters contained in the data files on the CD-ROM are:

Column Name	Units		
	. ,		
SITE_NAME	[none]		
SUB_SITE	[none]		
DATE_OBS	[DD-MON-YY]		
SOIL_DEPTH	[millimeters]		
RADON222_ACTIVITY	[becquerel][meter^-3]		
CRTFCN CODE	[none]		
REVISION DATE	[DD-MON-YY]		

7.3.4 Data Source

The sources of the parameter values contained in the data files on the CD-ROM are:

Column Name	Data Source		
SITE_NAME	[BORIS Designation]		
SUB_SITE	[BORIS Designation]		
DATE_OBS	Investigator		
SOIL_DEPTH	Investigator		
RADON222_ACTIVITY	Pylon alpha scintillation counters		
CRTFCN_CODE	[BORIS Designation]		
REVISION_DATE	[BORIS Designation]		

7.3.5 Data Range

Column Name

The following table gives information about the parameter values found in the data files on the CD-ROM.

Maximum

Data

Value

Missng Unrel Below

Data

Data

Value

Data

Detect Not

Value Limit Cllctd

Minimum

Data

Value

_	NSA-999-GMR02 TGB12-RDN01 15-NOV-93		None None None		None None None	None None None
_			None	None	None	None
RADON222 ACTIVITY						None
CRTFCN CODE		CPI				
REVISION DATE						
REVISION_DATE	21-A0G-90	03-1107-30	none	None	none	none
Minimum Data Value The minimum value found in the column. Maximum Data Value The maximum value found in the column. Missng Data Value The value that indicates missing data. This is used to indicate that an attempt was made to determine the parameter value, but the attempt was unsuccessful.						to
Unrel Data Value	to indicate a parameter val	The value that indicates unreliable data. This is used to indicate an attempt was made to determine the parameter value, but the value was deemed to be unreliable by the analysis personnel.				
Below Detect Limit	instruments de indicate that parameter val that the param	The value that indicates parameter values below the instruments detection limits. This is used to indicate that an attempt was made to determine the parameter value, but the analysis personnel determined that the parameter value was below the detection limit of the instrumentation.				
Data Not Cllctd	This value indetermine the indicates that not identical	dicates that no parameter valu t BORIS combine data sets into icular science	attempt e. This d severa the sam	usuall l simil e data :	Y ar but	le

Blank -- Indicates that blank spaces are used to denote that type of value. N/A -- Indicates that the value is not applicable to the respective column. None -- Indicates that no values of that sort were found in the column.

7.4 Sample Data Record

The following are wrapped versions of data record from a sample data file on the CD-ROM.

```
SITE_NAME, SUB_SITE, DATE_OBS, SOIL_DEPTH, RADON222_ACTIVITY, CRTFCN_CODE, REVISION_DATE

'NSA-OJP-FLXTR', 'TGB12-RDN03', 15-NOV-93, 40.0, 360, 'CPI', 21-AUG-96
'NSA-OJP-FLXTR', 'TGB12-RDN03', 15-NOV-93, 130.0, -999, 'CPI', 21-AUG-96
'NSA-OJP-FLXTR', 'TGB12-RDN03', 15-NOV-93, 240.0, -999, 'CPI', 21-AUG-96
'NSA-OJP-FLXTR', 'TGB12-RDN03', 15-NOV-93, 550.0, 5640, 'CPI', 21-AUG-96
'NSA-OJP-FLXTR', 'TGB12-RDN03', 15-NOV-93, 850.0, 7540, 'CPI', 21-AUG-96
'NSA-999-GMR02', 'TGB12-RDN01', 16-NOV-93, 20.0, 846, 'CPI', 21-AUG-96
'NSA-999-GMR02', 'TGB12-RDN01', 16-NOV-93, 85.0, 3570, 'CPI', 21-AUG-96
```

8. Data Organization

8.1 Data Granularity

The smallest unit of data is the 222Rn activity data for a given site on a given day.

8.2 Data Format(s)

The Compact Disk-Read-Only Memory (CD-ROM) files contain American Standard Code for Information Interchange (ASCII) numerical and character fields of varying length separated by commas. The character fields are enclosed with single apostrophe marks. There are no spaces between the fields.

Each data file on the CD-ROM has four header lines of Hyper-Text Markup Language (HTML) code at the top. When viewed with a Web browser, this code displays header information (data set title, location, date, acknowledgments, etc.) and a series of HTML links to associated data files and related data sets. Line 5 of each data file is a list of the column names, and line 6 and following lines contain the actual data.

9. Data Manipulations

9.1 Formulae

9.1.1 Derivation Techniques and Algorithms

Methods for calculating radon and radiocarbon data are given in Davidson and Trumbore (1995) and Trumbore (1995).

9.2 Data Processing Sequence

None given.

9.2.1 Processing Steps

None given.

9.2.2 Processing Changes

None given.

9.3 Calculations

None given.

9.3.1 Special Corrections/Adjustments

None given.

9.3.2 Calculated Variables

None given.

9.4 Graphs and Plots

None given.

10. Errors

10.1 Sources of Error

Sources of error include inaccuracies in soil collection and instrument errors during running of samples.

10.2 Quality Assessment

10.2.1 Data Validation by Source

There are many sources of error in flux measurements. The ones associated with the methods used here are detailed in Stephens and Sundquist (1998). Errors involved in soil gas sampling include the possibility that the volume of soil air space sampled (500 cc to 1 liter) represents a far larger volume of soil. Thus, the depth assigned to a soil gas sample (i.e., the depth of the probe) may not represent the average for the interval integrated in the gas sample taken from it (that is, air may be pulled down from above or up from below). Only analytical errors are given in these data sets.

10.2.2 Confidence Level/Accuracy Judgment

For radon measurements, accuracy is 10% of the measured value, with most of the error caused by uncertainty in the efficiency of the counting cell.

10.2.3 Measurement Error for Parameters

For radon measurements, accuracy is 10% of the measured value, with most of the error caused by uncertainty in the efficiency of the counting cell.

10.2.4 Additional Quality Assessments

None.

10.2.5 Data Verification by Data Center

Data were examined for general consistency and clarity.

11. Notes

11.1 Limitations of the Data

None given.

11.2 Known Problems with the Data

None given.

11.3 Usage Guidance

None given.

11.4 Other Relevant Information

None.

12. Application of the Data Set

The data can be used for estimation of soil gas exchange rates from ²²²Rn data (or testing of soil gas exchange rates derived from models).

13. Future Modifications and Plans

None given.

14. Software

14.1 Software Description

Calculations of fluxes for radon used linear regression from programs such as Microsoft Excel and Kaleidograph.

14.2 Software Access

None.

15. Data Access

The TGB-12 ²²²Rn activity data are available from the Earth Observing System Data and Information System (EOSDIS) Oak Ridge National Laboratory (ORNL) Distributed Active Archive Center (DAAC).

15.1 Contact Information

For BOREAS data and documentation please contact:

ORNL DAAC User Services Oak Ridge National Laboratory P.O. Box 2008 MS-6407 Oak Ridge, TN 37831-6407 Phone: (423) 241-3952

Phone: (423) 241-3952 Fax: (423) 574-4665

E-mail: ornldaac@ornl.gov or ornl@eos.nasa.gov

15.2 Data Center Identification

Earth Observing System Data and Information System (EOSDIS) Oak Ridge National Laboratory (ORNL) Distributed Active Archive Center (DAAC) for Biogeochemical Dynamics http://www-eosdis.ornl.gov/.

15.3 Procedures for Obtaining Data

Users may obtain data directly through the ORNL DAAC online search and order system [http://www-eosdis.ornl.gov/] and the anonymous FTP site [ftp://www-eosdis.ornl.gov/data/] or by contacting User Services by electronic mail, telephone, fax, letter, or personal visit using the contact information in Section 15.1.

15.4 Data Center Status/Plans

The ORNL DAAC is the primary source for BOREAS field measurement, image, GIS, and hardcopy data products. The BOREAS CD-ROM and data referenced or listed in inventories on the CD-ROM are available from the ORNL DAAC.

16. Output Products and Availability

16.1 Tape Products

None.

16.2 Film Products

None.

16.3 Other Products

These data are available on the BOREAS CD-ROM series.

17. References

17.1 Platform/Sensor/Instrument/Data Processing Documentation Not applicable.

17.2 Journal Articles and Study Reports

Davidson, E.A. and S.E. Trumbore. 1995. Gas diffusivity and production of CO₂ in deep soils of the eastern Amazon. Tellus. 47B: 550-565.

Mathieu, G.G. P.E. Biscaye, R.A. Lupton, and D.E. Hammond. 1988. System for measurement of ²²²Rn at low levels in natural waters. Heath Physics 55:989-992.

Newcomer, J., D. Landis, S. Conrad, S. Curd, K. Huemmrich, D. Knapp, A. Morrell, J. Nickeson, A. Papagno, D. Rinker, R. Strub, T. Twine, F. Hall, and P. Sellers, eds. 2000. Collected Data of The Boreal Ecosystem-Atmosphere Study. NASA. CD-ROM.

Sellers, P. and F. Hall. 1994. Boreal Ecosystem-Atmosphere Study: Experiment Plan. Version 1994-3.0, NASA BOREAS Report (EXPLAN 94).

Sellers, P. and F. Hall. 1996. Boreal Ecosystem-Atmosphere Study: Experiment Plan. Version 1996-2.0, NASA BOREAS Report (EXPLAN 96).

Sellers, P., F. Hall, and K.F. Huemmrich. 1996. Boreal Ecosystem-Atmosphere Study: 1994 Operations. NASA BOREAS Report (OPS DOC 94).

Sellers, P., F. Hall, and K.F. Huemmrich. 1997. Boreal Ecosystem-Atmosphere Study: 1996 Operations. NASA BOREAS Report (OPS DOC 96).

Sellers, P., F. Hall, H. Margolis, B. Kelly, D. Baldocchi, G. den Hartog, J. Cihlar, M.G. Ryan, B. Goodison, P. Crill, K.J. Ranson, D. Lettenmaier, and D.E. Wickland. 1995. The boreal ecosystem-atmosphere study (BOREAS): an overview and early results from the 1994 field year. Bulletin of the American Meteorological Society. 76(9):1549-1577.

Sellers, P.J., F.G. Hall, R.D. Kelly, A. Black, D. Baldocchi, J. Berry, M. Ryan, K.J. Ranson, P.M. Crill, D.P. Lettenmaier, H. Margolis, J. Cihlar, J. Newcomer, D. Fitzjarrald, P.G. Jarvis, S.T. Gower, D. Halliwell, D. Williams, B. Goodison, D.E. Wickland, and F.E. Guertin. 1997. BOREAS in 1997: Experiment Overview, Scientific Results and Future Directions. Journal of Geophysical Research 102(D24): 28,731-28,770.

Stephens, B.B. and E.T. Sundquist. 1998. Measurements of soil surface gas fluxes using closed chamber techniques, submitted for publication

Trumbore, S.E. 1995. Use of isotopes and tracers in the study of emission and consumption of trace gases in terrestrial environments. Chapter 9 in: Matson, P and Harriss, R, (eds.). Biogenic Trace Gases: Measuring Emissions from Soil and Water, Blackwell, Oxford, p. 291-326.

Winston, G.C., E.T. Sundquist, B.B. Stephens, and S.E. Trumbore. 1997. Winter CO₂ fluxes in a boreal forest. Journal of Geophysical Research 102(D24):28,795-28,804.

17.3 Archive/DBMS Usage Documentation None.

18. Glossary of Terms

None given.

19. List of Acronyms

AEAC - Albers Equal-Area Conic AFM AMS - Airborne Fluxes and Meteorology - Accelerator Mass Spectrometer ASCII - American Standard Code for Information Interchange AVHRR - Advanced Very High Resolution Radiometer BOREAS - BOReal Ecosystem-Atmosphere Study BORIS - BOREAS Information System CD-ROM - Compact Disk-Read-Only Memory - Distributed Active Archive Center DAAC EOS - Earth Observing System EOSDIS - EOS Data and Information System GIS - Geographic Information System GSFC - Goddard Space Flight Center HTML - Hyper-Text Markup Language IRGA - Infrared Gas Analyzer NAD83 - North American Datum of 1983 NASA - National Aeronautics and Space Administration NSA - Northern Study Area
OBS - Old Black Spruce
OJP - Old Jack Pine ORNL - Oak Ridge National Laboratory PANP - Prince Albert National Park - Time Domain Reflectometry TDR TGB TM SSA URL - Trace Gas Biogeochemistry - Thematic Mapper - Southern Study Area - Uniform Resource Locator USGS - United States Geological Survey YJP - Young Jack Pine

20. Document Information

20.1 Document Revision Date

Written: 14-Nov-1997 Last Updated: 08-Jul-1999

20.2 Document Review Date(s)

BORIS Review: 10-Jul-1998

Science Review:

20.3 Document ID

20.4 Citation

For soils data, the U.S. Geological Survey open file reports (see Section 17.2) should be cited. For interpretations, cite submitted Journal of Geophysical Research papers (see Section 17.2). Other citations should refer to the BOREAS Information System (BORIS) data set.

If using data from the BOREAS CD-ROM series, also reference the data as:

Trumbore, S., J. Harden, E. Sundquist, and E. Davidson, "Input, Accumulation and Turnover of Carbon in BOREAS NSA Soils (TGB-12)." In Collected Data of The Boreal Ecosystem-Atmosphere Study. Eds. J. Newcomer, D. Landis, S. Conrad, S. Curd, K. Huemmrich, D. Knapp, A. Morrell, J. Nickeson, A. Papagno, D. Rinker, R. Strub, T. Twine, F. Hall, and P. Sellers. CD-ROM. NASA, 2000.

Also, cite the BOREAS CD-ROM set as:

Newcomer, J., D. Landis, S. Conrad, S. Curd, K. Huemmrich, D. Knapp, A. Morrell, J. Nickeson, A. Papagno, D. Rinker, R. Strub, T. Twine, F. Hall, and P. Sellers, eds. Collected Data of The Boreal Ecosystem-Atmosphere Study. NASA. CD-ROM. NASA, 2000.

20.5 Document Curator

20.6 Document URL

REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE November 2000	3. REPORT TYPE AND DATES COVERED Technical Memorandum			
4. TITLE AND SUBTITLE Technical Report Series on the Boreas TGB-12 222Rn Acti	oreal Ecosystem-Atmosphere		5. FUNDING NUMBERS 923		
6. AUTHOR(S) Susan Trumbore, Eric Sundquist, and Greg Winston Forrest G. Hall and Sara K. Conrad, Editors					
7. PERFORMING ORGANIZATION NAMI Goddard Space Flight Center Greenbelt, Maryland 20771	E(S) AND ADDRESS (ES)		8. PEFORMING ORGANIZATION REPORT NUMBER 2000-03136-0		
9. SPONSORING / MONITORING AGE National Aeronautics and Space Washington, DC 20546-0001		(ES)	10. SPONSORING / MONITORING AGENCY REPORT NUMBER TM—2000–209891 Vol. 245		
11. SUPPLEMENTARY NOTES S. Trumbore: University of California, Irvine; E. Sundquist and G. Winston: U.S.G.S., Woods Hole, Massachusetts; S.K. Conrad: Raytheon ITSS					
12a. DISTRIBUTION / AVAILABILITY STA Unclassified—Unlimited Subject Category: 43 Report available from the NASA 7121 Standard Drive, Hanover,	A Center for AeroSpace Info	-	12b. DISTRIBUTION CODE		

13. ABSTRACT (Maximum 200 words)

The BOREAS TGB-12 team made measurements of soil carbon inventories, carbon concentration in soil gases, and rates of soil respiration at several sites to estimate the rates of carbon accumulation and turnover in each of the major vegetation types. Sampling strategies for soils were designed to take advantage of local fire chronosequences, so that the accumulation of carbon in regrowing mosses could be determined. All the data are used to (1) calculate the inventory of carbon and nitrogen in moss and mineral soil layers at NSA sites, (2) determine the rates of input and turnover (using both accumulation since the last stand-killing fire and radiocarbon data), and (3) link changes in soil respiration rate to shifts in the ¹⁴C content of soil CO₂ to determine the average 'age' respired CO₂. These ²²²Rn activity data were collected from 15-Nov-1993 to 16-Aug-1994 over the NSA sites. They are useful for determining the rate of gas exchange between soil and the overlying atmosphere. The data in this data set are stored in tabular ASCII files.

14. SUBJECT TERMS BOREAS, trace gas biog	15. NUMBER OF PAGES 13 16. PRICE CODE		
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT UL